

## 11A 脸识别

同种动物之间相互识别脸既是很高级的视觉认知，也是社会认知。对此，科学发掘了很多现象，也有一些机理研究，但理解有限。

初级视皮层 V1 继续投射到更高的区域，分为识别 where（物体空间位置）的背侧通路和识别 what（物体本征）的腹侧通路。

腹侧通路从 V1 到 V2、V4、及更远的区域，其可分辨的图像特征越来越复杂（Kobatake and Tanaka, 1994）。颞下皮层（inferotemporal cortex, IT）可以识别更复杂的图形，如：圆、方块、多刺圆、手等（Gross, Bender and Rocha-Miranda, 1969; Gross, Rocha-Miranda and Bender, 1972）。

普林斯顿大学心理系科学家在猴的 IT（Gross, Rocha-Miranda and Bender, 1972; Perret *et al.*, 1982; Desimone *et al.*, 1984）和颞上回（superior temporal sulcus, STS）（Bruce *et al.*, 1981）发现了识别脸的细胞，其中 IT 的脸识别细胞几乎都对脸特异反应，而对其他物体没有反应（Desimone *et al.*, 1984）。

人识别脸的能力强于黑猩猩和猴（Rosenfeld and van Hoesen, 1979; Parr, 2011）。黑猩猩、猴、绵羊、鸟类（鸡和鸽）、狗等动物也有脸识别细胞（Kendrick and Baldwin, 1987; Ryan and Lea, 1994; Kendrick *et al.*, 1996; Pascalis and Kelly, 2009）。绵羊不仅有识别绵羊脸的细胞，还有识别人脸、狗脸的细胞（Kendrick and Baldwin, 1987）。羊羔一到两个月认识母亲的脸（Kendrick *et al.*, 1998）。雌绵羊还对雄绵羊的脸有偏好（Kendrick *et al.*, 1995）。低等动物一般依赖嗅觉，但有两种蜂（wasps, *Polistes fuscatus* 和 *Polistes metricus*），*P fuscatus* 是群居的、*P metricus* 是独居的，前者有识别个体脸的能力，后者没有（Sheehan and Tibbets, 2011）。

用行为检测显示人对脸的关注在出生很早期就可能出现：9 分钟左右就对脸

的反应大于其他 (Goren, Sarty and Wu, 1975), 5 周就注视脸 (Haith, Bergman and Moore, 1977), 对脸是否好看也有不同的反应 (Slater, 1998, 2000)。在 4 天区分不带围巾的母亲与其他人的脸、35 天区分带围巾的母亲和其他人的脸 (Bushnell, Sai and Mullin, 1989; Walton, Bower and Bower, 1992; Pascalis *et al.*, 1995; Bruce *et al.*, 2000; Bartrip, Morton and De Schonen, 2001)。3 个月识别熟悉的脸 (De Haan *et al.*, 2001)。用 fMRI 检测观察到, 两个月的婴儿的对脸反应脑区被脸激活情况类似成人, 但脸还激活婴儿的语言区域 (Tzourio-Mazoyer *et al.*, 2002)。脸激活与成人一样在 9 岁儿童 (Gathers *et al.*, 2004)、或 12 岁 (Golarai *et al.*, 2007)。黑猩猩在 4 周左右识别母亲的脸 (Myowa-Yamakoshia, 2005)。

1980 年代的一系列电生理实验证明猴的神经元对脸有特异反应 (Bruce, Desimone and Gross, 1981; Perret, Rolls and Caan, 1982; Desimone *et al.*, 1984; Perrett *et al.*, 1984, 1985a, 1985, 1988b; Rolls, Baylis and Leonard, 1985; Saito *et al.*, 1986; Perret, Mistlin and Chitty, 1987)。STS 多感觉区域有只对脸反应的细胞 (Bruce *et al.*, 1981)。例如, 记录 497 个 STS 细胞, 48 个只对脸反应, 被脸持续激活, 28 个细胞在脸有转向、或颜色、大小、距离变化后反应不变 (Perret, Rolls and Caan, 1982)。早期在 IT 一次记录中, 41 个没有反应, 110 个有反应的细胞中, 66 个有选择性反应, 其中 20 对形状反应、2 个对手反应、3 个对脸有选择性反应 (Desimone *et al.*, 1984)。可以比较对脸和物体、脸和身体的反应, 找到对这三种分别有选择性反应的细胞 (Pinsk *et al.*, 2005)。通过 fMRI 辅助确定电生理电极插入位置, 可以找到特定区域内 97% 的细胞都对脸有选择性反应 (Tsao *et al.*, 2006; Fries, 2007; Fries, Tsao and Livingstone, 2007), 说明有脸特异区块 (patch)。用脑表面光学成像观察, 可以看到对脸呈现有选择性反应的脑区紧密

相连 (Wang, Tanaka and Tanifuji, 1996, 1998)。从而提出可能有脸朝向的功能柱 (Tanaka, 2003)。功能核磁共振实验也支持具有相似朝向选择性的面孔细胞在皮层上聚集 (Dubois *et al.*, 2015)。




图 11-18 倒置效应

脸识别能力有倒置效应，对正立的脸敏感性远远大于倒置的脸 (Yin, 1969; Thompson, 1980)。脸识别细胞对于脸的要求是一个圆加两点一杠 (大体相当于脸、眼和嘴) (Kobatake and Tanaka, 1994)。对脸有全面的识别和部件的敏感 (Freiwald, Tsao and Livingston, 2009)。在猴的脸识别细胞研究中，提出抑制性神经元可能对于脸识别很重要，去除 GABA 的抑制性作用后，原对脸 (和其他物体) 有特异反应的细胞失去反应特异性 (Wang, Fujita and Murayama, 2000)。用微电流刺激猴的面孔加工脑区 50-200 毫秒，可以增加其对脸的反应以及对个人面孔的识别 (Afraz, Kiani and Esteky, 2006; Moelle *et al.*, 2017)。

猕猴面孔脑区中较低级区域的神经元对面孔朝向非常敏感，而高级区域的神经元则可区分不同个体的面孔，且反应不依赖于面孔朝向，说明高级区域表征面

孔个体这一抽象概念 (Freiwald and Tsao, 2010)。进一步的研究对面孔个体的具体编码方式进行了探索,用计算模型生成上千张参数化的面孔,给动物呈现面孔图片的同时记录猕猴的面孔脑区,发现面孔细胞的反应和面孔模型中的抽象特征呈简单的线性关系,从~200 个细胞的反应可相当准确地重构呈现给动物的原始面孔 (Chang and Tsao, 2017)。

人对脸反应的脑区类似于猴 (Tsao *et al.*, 2003; Tsao, Moellet and Freiwald, 2008; Pinsk *et al.*, 2009; Srihasam *et al.*, 2012)。在开颅手术的病人经过允许能用颅内记录事件相关电位改变 (Allison *et al.*, 1999; McCarthy *et al.*, 1999, Puce, Allison and McCarthy, 1999), 记录到脸特异反应。也直接记录到神经细胞对脸反应 (Kreiman, Koch and Fried, 2000)。更多的是用正电子扫描 (PET) (Sergent, Ohta and MacDonald, 1992; Haxby *et al.*, 1994) 和 fMRI (Malach *et al.*, 1995; Puce *et al.*, 1996; Clark *et al.*, 1996; Kanwisher, McDermott and Chun, 1997; McCarthy *et al.*, 1997)。可以分别观察几个脑区 (FFA、OFA 和 fSTS) 对脸的部件和构型的敏感性 (Liu, Harris and Kanwisher, 2010)。跨颅磁刺激 (TMS) 是一种研究脑功能的方法 (Walsh and Cowey, 2000)。用 TMS 作用于特定脑区, 可以观察到脸反应的变化 (Pitcher *et al.*, 2007, 2008, 2009)。FFA 对脸的部件和构型都敏感, OFA 和 fSTS 只对真的脸部件反应、对其构型不反应 (Dzhelyova, Ellison and Atkinson, 2010)。

人类有不能识别脸的个体, 诊断为脸盲 (prosopagnosia, faceblind) (Bodamer, 1947), 分为先天型 (发育型) 和获得型。脸盲者可以识别其他物体, 而不能识别脸 (Farah, Levinson and Klein, 1995; Farah, 1996; Henke *et al.*, 1998; Nunn, Postma and Pearson, 2001; Duchaine and Nakayama, 2005; Duchaine *et al.*, 2006; Li

and Song, 2007; Riddoch *et al.*, 2008)。也有患者可以识别脸但不能识别其他物体 (Feinberg *et al.*, 1994; Moscovitch, Winocur and Behrmann, 1997; McMullen, Fisk and Phillips, 2000; Germine *et al.*, 2011)。对于脸盲的机理, 有多种解释, 有些脸盲可能确实是脸识别能力的特异变化 (Duchaine, 2006)。

后天获得的脸识别异常, 可以是病变或外伤 (Yin, 1970; Meadow *et al.*, 1974; Landis *et al.*, 1986; Barton *et al.*, 2002; Bouvier and Engel, 2006; Schiltz *et al.*, 2006; Steeves *et al.*, 2006)。右脑单侧外伤就可以导致脸盲。如果可以在脑成像观察到病变部位, 有助于了解参与脸识别的脑区 (Riddoch *et al.*, 2008)。这些可以与在正常人脑进行的核磁共振成像、外科手术人脑电生理记录相辅相成 (Kanwisher, McDermott and Chun, 1997; Tsao *et al.*, 2003; Barraclough and Perret, 2011)。

双生子研究显示脸识别能力有高度遗传性 (Polk *et al.*, 2007; Wilmer *et al.*, 2010; Zhu *et al.*, 2010)。德国一个大学的调查显示 2.47% 脸盲率 (Kennerknecht *et al.*, 2006, 2007), 中国香港大学的调查显示 1.88% 的脸盲率 (Kennerknecht, Ho and Wong, 2008)。遗传性脸盲在一些家系发现 (McConachie, 1976; Grueter *et al.*, 2007; Duchaine, Germine and Nakayama, 2007; Schmalzl, Palermo and Coltheart, 2008; Lee *et al.*, 2009)。这些表明经典遗传学和基因组学可以用于研究脸识别。

### 参考文献

- Afraz SR, Kiani R and Esteky H (2006) Microstimulation of inferotemporal cortex influences face categorization. *Nature* 422:692-695.
- Allison T, Puce A, Spencer DD and McCarthy G (1999) Electrophysiological studies of human face perception. I: Potentials generated in occipitotemporal cortex by

- face and nonface stimuli. *Cerebral Cortex* 9:415-430.
- Barraclough NE and Perrett DI (2011) from single cells to social perception. *Philosophical Transactions of the Royal Society of London B* 366:1739-1752.
- Bartrip J, Morton J and De Schonen S (2001) Responses to mother's face in 3-week to 5-month-old infants. *British Journal of Developmental Psychology* 19:219-232.
- Bodamer J (1947) Die prosopagnosie. *Archiv für Psychiatrie und Nervenkrankheiten* 179:6-53.
- Bruce CJ, Desimone R and Gross CG (1981) Visual properties of neurons in a polysensory area in superior temporal sulcus of the macaque. *Journal of Neurophysiology* 46:369-384.
- Bruce V, Campbell RN, Doherty-Sneddon G, Import A, Langton S, McAuley S and Wright R (2000) Testing face processing skills in children. *British Journal of Developmental Psychology* 18:319-333.
- Bushnell IW, Sai F and Mullin JT (1989) Neonatal recognition of the mother's face. *British Journal of Developmental Psychology* 7:3-15.
- Chang L and Tsao DY (2017) The code for facial identity in the primate brain. *Cell* 169:1013-1028.
- Clark VP, Keil K, Maisog JM, Courtney S, Ungerleider LG and Haxby JV (1996) Functional magnetic resonance imaging of human visual cortex during face matching: a comparison with positron emission tomography. *NeuroImage* 4:1-15.
- de Haan EH (1999) A familial factor in the development of face recognition deficits. *Journal of Clinical and Experimental Neuropsychology* 21: 312-315.

- de Haan M, Johnson MH, Maurer D and Perrett DI (2001) Recognition of individual faces and average face prototype by 1- and 3-month-old infants. *Cognitive Development* 16:1-20.
- Desimone R, Albright TD, Gross CG and Bruce CJ (1984) Stimulus-selective properties of inferior temporal neurons in the macaque. *Journal of Neuroscience* 4:2051-2062.
- Dubois J, de Berker AO and Tsao DY (2015) Single-unit recordings in the macaque face patch system reveal limitations of fMRI MVPA. *Journal of Neuroscience* 35:2791-2802.
- Duchaine B and Nakayama K (2005) Dissociations of face and object recognition in developmental prosopagnosia. *Journal of Cognitive Neuroscience* 17:249-261.
- Duchaine B, Yovel G, Butterworth E and Nakayama K (2006) Prosopagnosia as an impairment to face-specific mechanisms: Elimination of the alternative hypotheses in a developmental case. *Cognitive Neuropsychology* 23:714-747.
- Duchaine B, Germine L and Nakayama K (2007) Family resemblance: ten family members with prosopagnosia and within-class object agnosia. *Cognitive Neuropsychology* 24:419-430.
- Dzhelyova MP, Ellison A and Atkinson AP (2010) Event-related repetitive TMS reveals distinct, critical roles for right OFA and bilateral posterior STS in judging the trustworthiness and sex of faces. *Journal of Cognitive Neuroscience* 22:203-211.
- Farah MJ (1996) Is face recognition “special”? Evidence from neuropsychology.

*Behavioral Brain Research* 76:181-189.

Farah MJ, Levinson KL and Klein KL (1995) Face perception and within category discrimination in prosopagnosia. *Neuropsychologia* 33:661-674.

Feinberg TE, Schindler RJ, Ochoa E, Kwan PC and Farah MJ (1994) Associative visual agnosia and alexia without prosopagnosia. *Cortex* 30:395-412.

Freiwald WA, Tsao DY and Livingstone MS (2009) A face feature space in the macaque temporal lobe. *Nature Neuroscience* 12:1187-1196.

Freiwald WA and Tsao DY (2010) Functional compartmentalization and viewpoint generalization within the macaque face-processing system. *Science* 330:845-851.

Gathers AD, Bhatt R, Corbly CR, Farley AB and Joseph JE (2004) Developmental shifts in cortical loci for face and object recognition. *NeuroReport* 15:1549-1553.

Germine L, Cashdollar N, Duzel E and Duchaine B (2011) A new selective developmental deficit: Impaired object recognition with normal face recognition. *Cortex* 47:598-607.

Golarai G, Ghahremani DG, Whitfield-Gabrieli S, Reiss A, Eberhardt JL, Gabrieli JDE and Grill-Spector K (2007) Differential development of high-level cortex correlates with category-specific recognition memory. *Nature Neuroscience* 10:512-522.

Goren CC, Sarty M and Wu PYK (1975) Visual following and pattern discrimination of facelike stimuli by newborn infants. *Pediatrics* 56:544-549.

Gross CG, Bender DB and Rocha-Miranda CE (1969) Visual receptive fields of neurons in inferotemporal cortex of the monkey. *Science* 166:1303-1306.



- Gross CG, Rocha-Miranda CE and Bender DB (1972) Visual properties of neurons in inferotemporal cortex of the Macaque. *Journal of Neurophysiology* 35:96-111.
- Grueter M, Grueter T, Bell V, Halligan PW, Horst J, Sperling K, Halligan PW, Ellis HD and Kennerknecht I (2007) Hereditary prosopagnosia: the first case series. *Cortex* 43:734-749.
- Haith MM, Bergman TT and Moore MJ (1977) Eye contact and face scanning in early infancy. *Science* 198:853–855.
- Haxby JV, Horwitz B, Ungerleider LG, Maisog JM, Pietrini P and Grady CL (1994) The functional organization of human extrastriate cortex: a PET-rCBF study of selective attention to faces and locations. *Journal of Neuroscience* 14:6336-6353.
- Henke K, Schweinberger SR, Grigo A, Klos T and Sommer W (1998) Specificity of fact recognition: recognition of exemplars of non-face objects in prosopagnosia. *Cortex* 34:289-296.
- Kanwisher NG, McDermott J and Chun MM (1997) The fusiform face area: A module in human extrastriate cortex specialized for face perception. *Journal of Neuroscience* 17:4302-4311.
- Kendrick KM and Baldwin BA (1987) Cells in temporal cortex of conscious sheep can respond preferentially to the sight of faces *Science* 236:448-450.
- Kendrick KM, Atkins K, Hinton MR, Broad KD, Fabre-Nys C and Keverne B (1995) Facial and vocal discrimination in sheep. *Animal Behavior* 49:1665–1676.
- Kendrick KM, Atkins K, Hinton MR, Heavens P and Keverne B (1996) Are faces special for sheep? Evidence from facial and object discrimination learning tests

showing effects of inversion and social familiarity. *Behavioral Processes* 38:19-35.

Kennerknecht I, Grueter T, Welling B, Wentzek S, Horst J, Edwards S and Grueter M (2006) First report of prevalence of non-syndromic hereditary prosopagnosia (HPA). *American Journal of Medical Genetics* 140:1617-1622.

Kennerknecht I, Plümpe N, Edwards S and Raman R (2007) Hereditary prosopagnosia (HPA): the first report outside the Caucasian population. *Journal of Human Genetics* 52:230-236.

Kennerknecht I, Ho NY and Wong VC (2008) Prevalence of hereditary prosopagnosia (HPA) in Hong Kong Chinese population. *American Journal of Medical Genetics* 146A:2863-2870.

Kobatake E and Tanaka K (1994) Neuronal selectivities to complex object features in the ventral visual pathway of the macaque cerebral cortex. *Journal of Neurophysiology* 71:856–867.

Kreiman G, Koch C and Fried I (2000). Category specific visual responses of single neurons in the human medial temporal lobe. *Nature Neuroscience* 3:946-953.

Lee Y, Duchaine B, Wilson H and Nakayama K (2010) Three cases of developmental prosopagnosia from one family: detailed neuropsychological and psychophysical investigation of face processing. *Cortex* 46:949-964.

Li XB and Song YY (2007) First congenital progopagnosia case reported in China. *Progress in Natural Sciences* 17:165-171.

Liu J, Harris A and Kanwisher N (2010) Perception of face parts and face

configuration: an fMRI study. *Journal of Cognitive Neuroscience* 22:203-211.

Malach R, Reppas JB, Benson RB, Kwong KK, Jiang H, Kennedy WA, Ledden PJ, Brady TJ, Rosen BR and Tootell RBH (1995) Object-related activity revealed by functional magnetic resonance imaging in human occipital cortex. *Proceedings of the National Academy of Sciences USA* 92:8135-8138.

McCarthy G, Puce A, Gore J and Allison T (1997) Face-specific processing in the human fusiform gyrus. *Journal of Cognitive Neuroscience* 9:605-610.

McCarthy G, Puce A, Belger A and Allison T (1999) Electrophysiological studies of human face perception: II. Response properties of face-specific potentials generated in occipitotemporal cortex. *Cerebral Cortex* 9:431-444.

McConachie HR (1976) Developmental prosopagnosia. A single case report. *Cortex* 12: 76-82.

McMullen P, Fisk JD and Phillips S (2000) Apperceptive agnosia and face recognition. *Neurocase* 6:403-414.

Moeller S, Crapse T, Chang L and Tsao DY (2019) The effect of face patch microstimulation on perception of faces and objects. *Nature Neuroscience* 20:743-752.

Moscovitch M, Winocur G and Behrmann M (1997) What is special about face recognition? Nineteen experiments on a person with visual object agnosia and dyslexia but normal face recognition. *Journal of Cognitive Neuroscience* 9:555-604.

Myowa-Yamakoshia M, Yamaguchib MK, Tomonagac M, Tanakac M and

- Matsuzawa T (2005) Development of face recognition in infant chimpanzees (Pan troglodytes). *Cognitive Development* 20:49-63.
- Nunn JA, Postma P and Pearson R (2001) Developmental prosopagnosia: Should it be taken at face value? *Neurocase* 7:15-27.
- Parr LA (2011) The evolution of face processing in primates. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 366:1764-1777.
- Pascalis O, de Schonen S, Morton J, Deruelle C and Fabre-Grenet M (1995) Mother's face recognition by neonates: A replication and an extension. *Infant Behavioral Development* 18:79-85.
- Pascalis O and Kelly DJ (2009) The origins of face processing in humans: phylogeny and ontogeny. *Psychological Sciences* 4:200-209.
- Perret DI, Rolls ET and Caan W (1982) Visual neurons responsive to faces in the monkey temporal cortex. *Experimental Brain Research* 47:329-342.
- Perrett DI, Smith PAJ, Potter DD, Mistlin AJ and Head AS (1984) Neurons responsive to faces in the temporal cortex: Studies of functional organization, sensitivity to identity and relation to perception. *Human Neurobiology* 3:197-208.
- Perrett DI, Smith PAJ, Potter DD, Mistlin AJ, Head AS, Milner AD and Jeeves MA (1985a) Visual cells in the temporal cortex sensitive to face view and gaze direction. *Proceedings of the Royal Society of London Series B* 223:293-317.
- Perrett DI, Smith PAJ, Mistlin AJ, Chitt AJ, Head AS, Potter DD, Broennimann R, Milner AD and Jeeves MA (1985b) Visual analysis of body movement by

- neurones in the temporal cortex of the macaque monkey: a preliminary report. *Brain Research* 16:153-170.
- Perrett DI, Mistlin AJ and Chitty AJ (1987) Visual cells responsive to faces *Trends in Neuroscience* 10: 358-364.
- Perrett DI, Mistlin AJ, Chitty AJ, Smith PAJ, Potter DD, Broennimann R and Harries MH (1988) Specialized face processing and hemispheric asymmetry in man and monkey: Evidence from single unit and reaction time studies. *Behavioral Brain Research* 29:245-258.
- Pinsk MA, Desimone K, Moore T, Gross CG and Kastner S (2005) Representations of faces and body parts in macaque temporal cortex: A functional MRI study. *Proceedings of the National Academy of Sciences USA* 102:6996-7001.
- Pinsk MA, Arcaro M, Weiner KS, Kaljus JF, Inati SJ, Gross CG and Kastner S (2009) Neural representations of faces and body parts in macaque and human cortex: a comparative fMRI study. *Journal of Neurophysiology* 101:2581-2600.
- Pitcher D, Walsh V, Yovel G and Duchaine B (2007) TMS evidence for the involvement of the right occipital face area in early face processing. *Current Biology* 17:1566-1573.
- Pitcher D, Garrido L, Walsh V and Duchaine BC (2008) Transcranial magnetic stimulation disrupts the perception and embodiment of facial expressions. *Journal of Neuroscience* 28:8929-8933.
- Pitcher D, Charles L, Devlin JT, Walsh V and Duchaine B (2009) Triple dissociation of faces, bodies, and objects in extrastriate cortex. *Current Biology* 19:319-324.

- Polk TA, Park J, Smith MR and Park DC (2007) Nature versus nurture in ventral visual cortex: A functional magnetic resonance imaging study of twins. *Journal of Neuroscience* 27:13921-13925.
- Puce A, Allison T, Asgari M, Gore JC and McCarthy G (1996) Differential sensitivity of human visual cortex to faces, letterstrings, and textures: a functional magnetic resonance imaging study. *Journal of Neuroscience* 16:5205-5215.
- Puce A, Allison T and McCarthy G (1999) Electrophysiological studies of human face perception. III: Effects of top-down processing on face-specific potentials. *Cerebral Cortex* 9:445-458.
- Riddoch MJ, Johnston RA, Bracewell RM, Boutsen L and Humphreys GW (2008) Are faces special? A case of pure prosopagnosia. *Cognitive Neuropsychology* 25:3-26.
- Rolls ET (1984) Neurons in the cortex of the temporal lobe and in the amygdala of the monkey with responses selective for faces. *Human Neurobiology* 3:209-222.
- Rolls ET, Baylis GC and Leonard LM (1985) Role of low and high spatial frequencies in the face-selective responses of neurons in the cortex in the superior temporal sulcus in the monkey. *Vision Research* 25:1021-1035.
- Rosenfeld SA and Van Hoesen GW (1979) Face recognition in the rhesus monkey. *Neuropsychologia* 17:503-509.
- Ryan CME and Lea SEG (1994) Images of conspecifics as categories to be discriminated by pigeons and chickens: slides, video tapes, stuffed birds and live birds. *Behavioral Processes* 33:155-176.

- Saito H, Yukie M, Tanaka K, Hikosaka K, Fukada Y and Iwai E (1986) Integration of direction signals of image motion in the superior temporal sulcus of the macaque monkey. *Journal of Neuroscience* 6:145-157.
- Schmalzl L, Palermo R and Coltheart M (2008) Cognitive heterogeneity in genetically based prosopagnosia: A family study. *Journal of Neuropsychology* 2:99-117.
- Sergent J, Ohta S and MacDonald B (1992) Functional neuroanatomy of face and object processing: a positron emission tomography study. *Brain* 115:15-36.
- Sheehan MJ and Tibbets EA (2011) Specialized face learning is associated with individual recognition in paper wasps. *Science* 334:1272-1275.
- Slater A, von der Schulenburg C, Brown E, Badenoch M, Butterworth G, Parsons S and Samuels C (1998) Newborn infants prefer attractive faces. *Infant Behavioral Development* 21:345-354.
- Slater A, Bremner G, Johnson SP, Sherwood P, Hayes R, and Brown E (2000) Newborn infants' preferences for attractive faces: The role of internal and external facial features. *Infancy* 1:265-274.
- Srihasam K, Mandeville JB, Morocz IA, Sullivan KJ and Livingstone MS (2012) Behavioral and anatomical consequences of early versus late symbol training in macaques. *Neuron* 73:608-619.
- Tanaka K (2003) Columns for complex visual object features in the inferotemporal cortex: clustering of cells with similar but slightly different stimulus selectivities. *Cerebral Cortex* 13:90-99.
- Tate AJ, Fisher H, Leigh AE and Kendrick KM (2006) Behavioral and

neurophysiological evidence for face identity and face emotion processing in animals. *Philosophical Transactions of the Royal Society of London B Biological Sciences* 361:2155-2172.

Thompson P (1980) Margaret Thatcher: a new illusion. *Perception* 9:483-484.

Tsao DY, Freiwald WA, Knutsen TA, Mandeville JB and Tootell RB (2003) Faces and objects in macaque cerebral cortex. *Nature Neuroscience* 6:989-95.

Tsao DY, Freiwald WA, Tootell RB and Livingstone MS (2006) A cortical region consisting entirely of face selective cells. *Science* 311:670-674.

Tsao DY, Moellet S and Freiwald WA (2008) comparing face patch systems in macaques and humans. *Proceedings of the National Academy of Sciences USA* 105:19514-19519.

Tzourio-Mazoyer N, de Schonen S, Crivello F, Reutter B, Aujard Y and Mazoyer BM (2002) Neural correlates of woman face processing by 2-month-old infants. *NeuroImage* 15:454-461.

Walsh V and Cowey A (2000) transcranial magnetic stimulation and cognitive neuroscience. *Nature Reviews Neuroscience* 1:73-79.

Walton GE, Bower NJA and Bower TGR (1992) Recognition of familiar faces by newborns. *Infant Behavioral Development* 15:265-269.

Wang G, Tanaka K and Tanifuji M (1996) Optical imaging of functional organization in the monkey inferotemporal cortex. *Science* 272:1665-1668.

Wang G, Tanifuji M and Tanaka K (1998) Functional architecture in monkey inferotemporal cortex revealed by *in vivo* optical imaging. *Neuroscience*



*Research* 32:33-46.

Wilmer JB, Germine L, Chabris CF, Chatterjee G, Williams M, Loken E, Nakayama K and Duchaine B (2010). Human face recognition ability is specific and highly heritable. *Proceedings of the National Academy of Sciences USA* 107: 5238–5241.

Yin RK (1969) Looking at upside-down faces. *Journal of Experimental Psychology* 81:141-145.

Zhu Q, Song Y, Hu S, Li X, Tian M, Zhen Z, Dong Q, Kanwisher N and Liu J (2010) Heritability of the specific cognitive ability of face perception. *Current Biology* 20:137-142.